

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B16

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61L 27/22, 27/38, A61K 38/18, C07K 14/51		A1	(11) International Publication Number: WO 00/44413 (43) International Publication Date: 3 August 2000 (03.08.00)
(21) International Application Number: PCT/US00/02430 (22) International Filing Date: 31 January 2000 (31.01.00) (30) Priority Data: 60/118,160 1 February 1999 (01.02.99) US 09/493,543 28 January 2000 (28.01.00) US (71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 Cambridge Park Drive, Cambridge, MA 02140 (US). (72) Inventors: ZHANG, Renwen; 10 Matthew Court, Edison, NJ 08820 (US). PELUSO, Diane; 107 Emory Road, Marshfield, MA 02050 (US). MORRIS, Elisabeth; 15 Apple Street, Sherborn, MA 01770 (US). (74) Agent: GYURE, Barbara, A.; American Home Products Corporation, Patent and Trademark Department-2B, One Campus Drive, Parsippany, NJ 07054 (US).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS AND COMPOSITIONS FOR HEALING AND REPAIR OF ARTICULAR CARTILAGE			
(57) Abstract Methods and compositions are provided for the treatment of articular cartilage defects and disease involving the combination of the tissue, such as osteochondral grafts, with active growth factor. The active growth factor is preferably a composition containing at least one bone morphogenetic protein and a suitable carrier. The method results in the regeneration of functional repair of articular cartilage tissue.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS AND COMPOSITIONS FOR HEALING AND REPAIR OF ARTICULAR CARTILAGE

FIELD OF THE INVENTION

5 The present invention relates to the field of tissue repair, specifically, the regeneration of stable and functional articular cartilage repair. Thus, the present invention may be useful in reconstructive surgery or other procedures for the regeneration or repair of articular cartilage.

BACKGROUND OF THE INVENTION

10 The repair of articular cartilage injuries remains a challenge in present day orthopedics. Several of the current therapeutic strategies are based upon the grafting of chondral and osteochondral tissues. Autologous osteochondral grafting provides the most appropriate physiological material. However, donor tissue is limited, and often requires surgery at a secondary site in order to harvest tissue for transplant. Accordingly, despite
15 substantial endeavors in this field, there remains a need for an effective method of repair of articular cartilage defects and injuries which provides appropriate physiological repair without the need to collect autologous tissue from the patient.

SUMMARY OF THE INVENTION

20 The present invention provides methods and compositions for regenerating functional and physiologically appropriate tissue repair for the repair of articular cartilage injuries and defects. In particular, the present invention comprises methods of treating patients with articular cartilage injuries or defects. The methods and compositions of the present invention are advantageous in that they utilize bone morphogenetic proteins (BMPs), which are known to have osteogenic and/or chondrogenic properties, and which
25 may be produced via recombinant DNA technology, and therefore are of potentially unlimited supply. The methods and compositions of the present invention are further advantageous in that regeneration of functional articular cartilage may be accelerated or may be of greater ultimate strength and stability, and the tissue formed at the site of the

5 defect or injury is physiologically appropriate.

The use of BMP to augment the repair of articular cartilage defects and injuries may result in better methods for treatment of osteoarthritis, thus obviating, delaying or reducing the need for artificial hip replacements and other common interventions. Preclinical evaluations indicate that rhBMP-2 improves early healing of full thickness defects of articular cartilage in rabbits.

10

DETAILED DESCRIPTION OF THE INVENTION

According to the present invention, methods and compositions are provided for treatment of patients who suffer from some form of articular cartilage injury or defect. The injury may be the result of acute stress, or injury, such as resulting from participation in athletics, or from accidental occurrences which tear, mar or otherwise injure the articular cartilage.

15

The methods and composition are advantageous in that repair or improvement of articular cartilage defects, particularly full thickness articular cartilage defects. Other defects may also be treated by the methods and compositions of the present invention, particularly with an additional procedure in which the site of the defect is further aggravated so as to reach the underlying subchondral bone.

20

In the present invention, active growth factor, such as a BMP, is added to a suitable tissue source. The tissue source may be an osteochondral graft, either autologous to the patient, or may comprise allograft or artificially prepared tissue. In a preferred embodiment, the tissue source may be chondrocytic cell cultures, such as chondrocyte or stem cell cultures which have been prepared through *ex vivo* cell culture methods, with or without additional growth factors. For example, see the disclosure of US5226914; US5811094; US5053050; US5486359; US5786217 and US5723331. The disclosures of all of these applications are hereby incorporated herein by reference.

25

The tissue may also be harvested by traditional non-cell culture based means, using techniques such as mosaicplasty, in which cartilage is harvested using commercially available instruments such as Acufex7 [Smith and Nephew, Inc., Andover MA]; COR

30

- 5 System [Innovasive Technologies, Marlborough MA]; or Arthrex⁷ Osteochondral Autograft Transfer System [Arthrex, Munich, Germany]. The tissue harvested may be applied directly in the methods of the present invention, or may be combined with the tissue based cell culture systems described above.

GROWTH FACTOR

- 10 The active growth factor used in the present invention is preferably from the subclass of proteins known generally as bone morphogenetic proteins (BMPs), which have been disclosed to have osteogenic, chondrogenic and other growth and differentiation type activities. These BMPs include rhBMP-2, rhBMP-3, rhBMP-4 (also referred to as rhBMP-2B), rhBMP-5, rhBMP-6, rhBMP-7 (rhOP-1), rhBMP-8, rhBMP-9,
15 rhBMP-12, rhBMP-13, rhBMP-15, rhBMP-16, rhBMP-17, rhBMP-18, rhGDF-1, rhGDF-3, rhGDF-5, rhGDF-6, rhGDF-7, rhGDF-8, rhGDF-9, rhGDF-10, rhGDF-11, rhGDF-12, rhGDF-14. For example, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6 and BMP-7, disclosed in United States Patents 5,108,922; 5,013,649; 5,116,738; 5,106,748; 5,187,076; and 5,141,905; BMP-8, disclosed in PCT publication WO91/18098; and BMP-9,
20 disclosed in PCT publication WO93/00432, BMP-10, disclosed in United States Patent 5,637,480; BMP-11, disclosed in United States Patent 5,639,638, or BMP-12 or BMP-13, disclosed in United States Patent 5,658,882, BMP-15, disclosed United States Patent 5,635,372 and BMP-16, disclosed in co-pending patent application serial number 08/715,202. Other compositions which may also be useful include Vgr-2, and any of the
25 growth and differentiation factors [GDFs], including those described in PCT applications WO94/15965; WO94/15949; WO95/01801; WO95/01802; WO94/21681; WO94/15966; WO95/10539; WO96/01845; WO96/02559 and others. Also useful in the present invention may be BIP, disclosed in WO94/01557; HP00269, disclosed in JP Publication number: 7-250688; and MP52, disclosed in PCT application WO93/16099. The
30 disclosures of all of these applications are hereby incorporated herein by reference. Also useful in the present invention are heterodimers of the above and modified proteins or partial deletion products thereof. These proteins can be used individually or in mixtures

5 of two or more, and rhBMP-2 is preferred.

The BMP may be recombinantly produced, or purified from a protein composition. The BMP may be homodimeric, or may be heterodimeric with other BMPs (e.g., a heterodimer composed of one monomer each of BMP-2 and BMP-6) or with other members of the TGF- β superfamily, such as activins, inhibins and TGF- β 1 (e.g., a
10 heterodimer composed of one monomer each of a BMP and a related member of the TGF- β superfamily). Examples of such heterodimeric proteins are described for example in Published PCT Patent Application WO 93/09229, the specification of which is hereby incorporated herein by reference. The amount of osteogenic protein useful herein is that amount effective to stimulate increased osteogenic activity of infiltrating progenitor cells,
15 and will depend upon the size and nature of the defect being treated, as well as the carrier being employed. Generally, the amount of protein to be delivered is in a range of from about 0.05 to about 1.5 mg.

In a preferred embodiment, the osteogenic protein is administered together with an effective amount of a protein which is able to induce the formation of tendon- or
20 ligament-like tissue. Such proteins, include BMP-12, BMP-13, and other members of the BMP-12 subfamily, as well as MP52. These proteins and their use for regeneration of tendon and ligament-like tissue are disclosed in United States application serial number serial number 08/362,670, filed on December 22, 1994, the disclosure of which is hereby incorporated herein by reference. In another preferred embodiment, a heterodimer in
25 which one monomer unit is an osteogenic protein such as BMP-2, and the other monomer subunit is a tendon-inducing protein, such as BMP-12, is administered in accordance with the methods described below, in order to induce the formation of a functional attachment between connective tissue and bone.

5

APPLICATION OF GROWTH FACTOR

Growth factor may be applied to the tissue source in the form of a buffer solution. One preferred buffer solution is a composition comprising, in addition to the active growth factor, about 1.0 to about 10.0% (w/v) glycine, about 0.1 to about 5.0% (w/v) of a sugar, preferably sucrose, about 1 to about 20 mM glutamic acid hydrochloride, and optionally about 0.01 to about 0.1% of a non-ionic surfactant, such as polysorbate 80. Preferred solutions are from about 1% to about 20% w/v cellulosic carrier/buffer. If desired, a salt may be added.

Other materials which may be suitable for use in application of the growth factors in the methods and compositions of the present invention include hyaluronic acid, surgical mesh or sutures, polyglyconate, temperature-sensitive polymers, demineralized bone, minerals and ceramics, such as calcium phosphates, hydroxyapatite, etc., as well as combinations of the above described materials. In the preferred embodiment of the present invention, however, no carrier is employed.

The growth factor of the present invention, in a suitable buffer such as that described above, or combined with a suitable carrier, may be applied directly to the tissue and/or to the site in need of tissue repair. For example, the growth factor may be physically applied to the tissue through spraying or dipping, or using a brush or other suitable applicator, such as a syringe for injection. Alternatively, or in conjunction, the protein may be directly applied to the site in need of tissue repair.

The following examples further describe the practice of embodiments of the invention with BMP-2. The examples are not limiting, and as will be appreciated by those skilled in the art, can be varied in accordance with the above specification.

5

EXAMPLES

I. Rabbit Allograft

All procedures were carried out with approval from IACUC. Twelve male New Zealand white rabbits (6 months old) were used. Two rabbits served as donors and 10 as recipients. Osteochondral grafts (3.5 mm diameter) were harvested from the trochlear groove or the medial femoral condyle of the donors, and transplanted into a 3.5 mm deep defect in the trochlear groove of the recipient. The graft was bathed in either rhBMP-2 (0.5 mg/ml) or buffer control prior to implantation. The rabbits were sacrificed 4 weeks after surgery and the transplants and surrounding tissue were evaluated by a histologic-histochemical grading scale, as described in Sellers et al., *J. Bone Joint Surg.*, 79-A:1452-1463 (1997). Computerized image analysis of histologic sections was also performed. Results were evaluated using the unpaired Students t-test.

On gross examination, the joints showed no signs of inflammation. All the defects were filled by repair tissue. The surface appearance of the defects was variable but acceptable and did not correlate with form of treatment. Osteophytes were found in 3 joints (2 in the experimental group; 1 in control buffer group).

There was no correlation between the gross and histologic appearance in any of the defects. The presence of chondrocytes in the lacunae and sporadic cloning of cells in the donor cartilage indicated survival of the tissue. Focal degeneration of the donor cartilage was present in all of the control groups, but only one of the rhBMP-2 treated group. The healing of the defect in the rhBMP-2 treated group was significantly improved compared to that in the control group. The rhBMP-2 treated group had improved bony integration indicated by less fibrous repair tissue in the subchondral bone compartment. Treatment with rhBMP-2 also resulted in more cartilage above the original tidemark, apparently consisting of both donor tissue and newly regenerated recipient cartilage. There was no significant difference in the total amount of bone observed between the two groups.

5

TABLE I
HISTOLOGIC SCORE AND HISTOMORPHOMETRIC MEASUREMENT
FOR CARTILAGE REPAIR, MEAN VALUE (SD)

Parameter	rhBMP-2	Control
Average Score**	10.0 (5.42)*	20.6 (5.18)
% of bone under tidemark	73.26 (13.28)	62.88 (18.07)
% of fibrous tissue under tidemark	2.19 (2.04)	15.81 (9.88)
% of cartilage above tidemark	74.70 (41.08)*	18.17 (26.70)
% of filling of the defect	96.53 (4.86)*	88.79 (8.04)

15

* Statistically significant difference from control ($p < 0.05$).

** Scale system ranges from 0 (normal cartilage) to 31 (no repair).

Additional histomorphometric analysis data further supports the beneficial effects of rhBMP-2 on the healing of graft. For example, the percentage filling of the new tissue above tide marker has been shown to be 81.52% in a rhBMP-2 treated group vs. 57.63% in control. There was less graft cartilage degeneration in rhBMP-2 treated group (23.83%) than in control group (44.52%). The integration of the graft or newly formed cartilage with the host cartilage was improved by rhBMP-2 treatment (56.48%) compared to that of control group (21.89%). More new cartilage formed under the influence of rhBMP-2 either at the edge of graft, which eliminated the gap between the graft and host, or at the top of graft, which made the graft more congruent with the joint surface.

The above results demonstrate that the healing of allogeneic osteochondral grafts in articular cartilage defects was improved by the addition of rhBMP-2. The active growth factor may have accelerated subchondral bone union, providing support and nutrition to the articular cartilage tissue. Addition of growth factor may also have stimulated new cartilage formation from recipient mesenchymal stem cells in the bone marrow and/or the synovial tissue. These results suggest that the combination of active growth factor, particularly the bone morphogenetic proteins, and osteochondral allografts

- 5 might present a potent strategy for treatment of articular cartilage defects, particularly full thickness articular cartilage defects.

II. Rabbit Autograft

- 10 Osteochondral grafts (2.7 mm in diameter and 3.0 mm long) were harvested from the trochlear groove or femoral condyle and transplanted into a donor site 2.7mm wide and 3.5mm long on the trochlear groove or femoral condyle of the knee joint in rabbits. Half the animals had buffer dripped into the recipient site prior to transplantation, and then the grafts were dipped in buffer for 2 minutes and placed into the recipient site. The other half had 5µg rhBMP-2 dripped into the recipient site prior to transplantation, and then the graft was dipped into buffer containing 500µg/ml rhBMP-2 for 2 minutes and
15 then transplanted into the recipient site. The animals were sacrificed 4 weeks after surgery, and the recipient sites were evaluated histologically using both a histologic-histochemical grading scale [Sellers, et al., J. Bone Joint Surg., 79-A: 1452-63 (1997)] and quantitative computerized image analysis of the tissue. The data indicated that treatment with rhBMP-2 improved the healing of the autograft. The most dramatic effects
20 were the reduction of graft cartilage degeneration (rhBMP-2 8.18% vs. control 36.25%), and more cartilage formed at the edge of graft (rhBMP-2 88.23% vs. control 50%).

III. Non-Human Primate Autograft:

- 25 The non human primates used for autografts experiments were cynomologous macaques. Osteochondral grafts (3.5mm diameter x 6mm long) were harvested from the trochlear groove of 6 cynomologous macaques and transplanted into recipient sites drilled into both the medial and lateral femoral condyle of the same animal (n=12 transplants total). Prior to transplantation 25µg rhBMP-2 was dripped into 6 recipient sites, and the grafts from those 6 transplants were dipped into a solution of 1.25mg/ml rhBMP-2 for 2 minutes. In the other 6 transplants, buffer alone was dripped into the recipient sites and
30 the grafts were dipped into buffer alone for 2 minutes prior to transplantation. The limbs were immobilized in a cast for 2 weeks post-operatively, and the animals were sacrificed 9 weeks post operatively.

5 All the animals had normal function of their knee joints. On gross examination, the joints showed no signs of inflammation. Osteophytes were not found in any joint. Although the surface of the defects appeared level with the surrounding cartilage on gross examination, microscopic observation revealed subsidence of the grafts in most of the cases. The tissue observed grossly covering the surface was actually new-formed tissue
10 on the top of graft. Computerized image analysis was performed by a blinded evaluator to quantitate percent filling of the defect, the new tissue types formed above the original tide mark, and the integration of the grafts and the surrounding cartilage. Favorable results were observed in the rhBMP-2 treated group in all these parameters. More new cartilage formed between the graft and host cartilage to eliminate the gap resulting in better
15 integration of the graft with the surrounding cartilage (rhBMP-2 88.59% vs. control 64.82%). The filling of the cartilage defect was better in rhBMP-2 treated group (95.02%) than in the control group (86.68%). There was more fibrous tissue in the control group (11.90% vs. rhBMP-2 5.65%), while more transitional tissue was found in the rhBMP-2 treated group (36.38% vs. control 20.53%). There was no significant difference on the
20 overall histologic-histochemical score between the two groups. Peripheral quantitative computerized tomography (pQCT) showed that the bone density increased in the donor sites with time. At 6 weeks and 9 weeks after the operation, the tissue in the rhBMP-2 treated donor sites was significantly denser and the healing process was more advanced compared to control sites. Histologically, the donor sites contained regenerated bone trabeculae
25 with fibrous tissue at the surface in all the cases.

IV. rhBMP-2 Retention Ex Vivo:

Retention of rhBMP-2 in osteochondral graft with this technique was evaluated with the grafts from non-human primates. The graft was dipped in a mixture solution of ¹²⁵I labeled rhBMP-2 and unlabeled rhBMP-2. Results showed that the amount of rhBMP-
30 2 absorbed to graft was proportional to the concentration of the protein, and the time of soaking. Other factors, which affect the retention of rhBMP-2, included the size of graft, and the presence of marrow elements between trabecular bone.

5 **V. rhBMP-2 Retention Time Course In Vivo:**

10 The time course of rhBMP-2 retention in osteochondral graft was evaluated in rabbits. A mixture solution of ¹²⁵I labeled rhBMP-2 and unlabeled rhBMP-2, which contained 5 ug rhBMP-2 and 20 uCi ¹²⁵I, was loaded to the graft before implantation. The animals were scanned with γ-camera during the follow-up time for 22 days post-operatively. Compared to the time course of collagen sponge as a carrier, the half time of rhBMP-2 in osteochondral graft was increased from 1 day to 3 days. The radioactivity of 10% of the starting point was maintained from 11 days of collagen sponge to 22 days of graft.

15 **VI. Non-Human Primate Allografts:**

20 Donor sites (3.5 mm wide x 6 mm long) were removed from the trochlear grooves of 12 adult cynomolgous macaques and transplanted into 3.5 x 6mm recipient sites in the medial and lateral femoral condyles of unrelated individuals. Half of the transplants were soaked in 1.25 mg/ml rhBMP-2 for 2 minutes prior to transplantation, and half were soaked in buffer. The identical procedure was performed on the other limb 7 weeks after the first surgery. The limb was immobilized in a cast for 2 weeks post operatively after each surgery, and the animals were sacrificed 9 weeks after the second surgery for histologic analysis.

25 These results suggest that the combination of active growth factor, particularly the bone morphogenetic proteins, and osteochondral autografts might present a potent strategy for treatment of articular cartilage defects, particularly full thickness articular cartilage defects. In other embodiments BMP-2 may also be applied to frozen osteochondral allograft for treatment of focal articular cartilage defect.

30 The foregoing descriptions detail presently preferred embodiments of the present invention. Numerous modifications and variations in practice thereof are expected to occur to those skilled in the art upon consideration of these descriptions. Those modifications and variations are believed to be encompassed within the claims appended hereto.

5

CLAIMS**We claim:**

1. A method for regeneration of articular cartilage comprising administering to an area in need of regeneration of said articular cartilage at least one purified bone morphogenetic protein (BMP).
2. A method for regeneration of articular cartilage comprising administering to an area in need of regeneration of said articular cartilage a suitable tissue source in combination with at least one purified bone morphogenetic protein (BMP).
3. The method of claim 1 wherein said BMP is BMP-2.
4. The method of claim 2 wherein said BMP is BMP-2.
5. A method for regeneration of articular cartilage comprising administering to an area in need of regeneration of said articular cartilage at least one purified protein selected from the group consisting of Vgr-2, growth and differentiation factors (GDFs), and BIP.
6. The method of claim 1 further comprising a protein which induces the formation of tendon or ligament-like tissue.
7. The method of claim 6 wherein said protein which induces the formation of tendon or ligament-like tissue is selected from the group consisting of BMP-12, BMP-13 members of the BMP-12 subfamily and MP52.
8. A composition for regeneration of articular cartilage said composition comprising at least one purified bone morphogenetic protein (BMP).

- 5 9. A composition for regeneration of articular cartilage said composition comprising a suitable tissue source in combination with at least one purified bone morphogenetic protein (BMP).
- 10 10. The composition of claim 8 wherein said BMP is BMP-2
11. The composition of claim 9 wherein said BMP is BMP-2.
- 15 12. A composition for regeneration of articular cartilage said composition comprising at least one purified protein selected from the group consisting of Vgr-2, growth and differentiation factors (GDFs), and BIP.
13. The composition of claim 8 further comprising a protein which induces the formation of tendon or ligament-like tissue.
- 20 14. The composition of claim 13 wherein said protein which induces the formation of tendon or ligament-like tissue is selected from the group consisting of BMP-12, BMP-13 members of the BMP-12 subfamily and MP52.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/02430

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61L27/22 A61L27/38 A61K38/18 C07K14/51		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61L A61K C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 31788 A (GENETICS INST) 23 July 1998 (1998-07-23) page 4, line 9 - line 29; examples	1-14
X	TAKAGI K ET AL: "THE ROLE OF BONE MARROW IN BONE MORPHOGENETIC PROTEIN-INDUCED REPAIR OF FEMORAL MASSIVE DIAPHYSEAL DEFECTS" CLINICAL ORTHOPAEDICS AND RELATED RESEARCH, US, PHILADELPHIA, PA, vol. 171, 1 January 1982 (1982-01-01), pages 224-231, XP002065500 page 230	1-14
X	US 5 786 217 A (TUBO ROSS A ET AL) 28 July 1998 (1998-07-28) cited in the application claims; examples	1-4, 8-11
-/-		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"8" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">17 May 2000</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">07/06/2000</div>
Name and mailing address of the ISA European Patent Office, P.B. 6818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Facs (+31-70) 340-3018		Authorized officer <div style="text-align: center; font-weight: bold;">ESPINOSA, M</div>

INTERNATIONAL SEARCH REPORT

Inter national Application No
PCT/US 00/02430

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WANG E A: "BONE MORPHOGENETIC PROTEINS (BMPs): THERAPEUTIC POTENTIAL IN HEALING BONY DEFECTS" TRENDS IN BIOTECHNOLOGY, GB, ELSEVIER PUBLICATIONS, CAMBRIDGE, vol. 11, no. 9, 1 September 1993 (1993-09-01), pages 379-383, XP002065499 ISSN: 0167-7799 page 381 -page 382</p>	1-4,8-11
X	<p>REDDI A H ET AL: "BONE MORPHOGENETIC PROTEINS. POTENTIAL ROLE IN OSTEOPOROSIS" OSTEOPOROSIS, GB, SHEFFIELD, 1 January 1996 (1996-01-01), pages 281-287, XP002065501 ISSN: 1365-3326 page 285</p>	1,3,5-8, 10,12-14
X	<p>US 5 658 882 A (CELESTE ANTHONY J ET AL) 19 August 1997 (1997-08-19) cited in the application column 11, line 43 -column 67; claims</p>	1,3,6-8, 10,13,14
X	<p>WO 96 39170 A (GENETICS INST) 12 December 1996 (1996-12-12) claims; examples</p>	1,3,6-8, 10,13,14
X	<p>WO 94 01557 A (SUMITOMO METAL IND ;KANGAWA KENJI (JP); HINO JUN (JP); FUKUDA KENJ) 20 January 1994 (1994-01-20) cited in the application claims</p>	5,12

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter national Application No

PCT/US 00/02430

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9831788	A	23-07-1998	AU 6245898 A	07-08-1998
US 5786217	A	28-07-1998	AU 2435495 A	29-11-1995
			WO 9530742 A	16-11-1995
			US 5723331 A	03-03-1998
US 5658882	A	19-08-1997	US 6027919 A	22-02-2000
			AU 689184 B	26-03-1998
			AU 1301395 A	27-06-1995
			CA 2176942 A	15-06-1995
			EP 0733109 A	25-09-1996
			FI 962350 A	16-07-1996
			JP 9506261 T	24-06-1997
			NO 962304 A	04-06-1996
			WO 9516035 A	15-06-1995
WO 9639170	A	12-12-1996	US 5902785 A	11-05-1999
			AU 6024696 A	24-12-1996
WO 9401557	A	20-01-1994	AU 4514193 A	31-01-1994
			JP 6172390 A	21-06-1994

THIS PAGE BLANK (USPTO)